



<p>Uses</p>	<p>ASTYFER is a superior blood builder that is used to treat a person with iron deficiency anemia (lack of red blood cell). ASTYFER has ingredients that offers a faster blood cell formation. Where there is lack of red blood cell in the body, ASTYFER is the solution.</p> <p>How to use ASTYFER: Follow your doctor's prescription and if available, read the Patient Information Leaflet provided in the pack before you start taking ASTYFER capsule or syrup. If you have any questions, ask your doctor or pharmacist.</p> <p>Take ASTYFER capsule and syrup by mouth before or after meal as directed by your doctor or pharmacist.</p>
<p>How to Use</p>	

Side Effects	<p>Stomach upsets, feeling sick, <u>vomiting</u>, hard or difficult to poo, frequently watery poo, <u>black or darker than normal appearing stools</u>, high nitrogen (a colourless element that has no smell and is usually found as a gas) in blood.</p> <p>If any of these effects last or get worse, tell your doctor or pharmacist immediately. Remember that your doctor has prescribed this medication because he or she has judged that the benefit to you is greater than the risk of side effects.</p>
Precautions	<p>Before taking ASTYFER, inform your doctor or pharmacist about the current drugs you bought at the pharmacy (e.g. vitamins, herbal supplements, etc.), about your allergies to capsule or syrup, about pre-existing diseases, and about your current health condition (e.g pregnancy, upcoming surgery, etc.). The essence of this is to make sure that the drug is safe for you to use.</p>
Interactions	<p>Keep a list of all the products you use (including prescription/non-prescription drugs and herbal drugs) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.</p>

Dosage	<p>ASTYFER should be taken as your doctor or pharmacist prescribes. For children, the dosage is based on weight and medical condition.</p> <p>Do not take this medication more often or increase your dose without consulting your doctor. Your condition will not improve any faster and the risk of serious side effects may increase.</p> <p>Missed dose: If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up. “If symptoms persist or gets worse after 3 days, consult your doctor”</p>
Storage	<p>Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets.</p>